


Lindgren, CC + Ralph R Mellan, Nuclear Phenomena suggesting  
a sexual mechanism for the tubule bristles.  
Proc Soc 30:110 - 1932.


Mellan et al Proc Soc 30:80 1932

AFB  $\rightarrow$  filamentous gonidia  $\rightarrow$  var AF diplococci  $\rightarrow$   
tetrad diplococcus  $\rightarrow$  diploths  $\rightarrow$  actast gonidia  $\rightarrow$  R + bc  
 $\rightarrow$  S + bc

Acetocarmine pupus.



Tetranomus



Mauhiel, JG Contribution à l'étude de la variation en micologie.  
logie. Th. doc. de nat., Nancy 307p. 1932.  
: from ~~Annales~~ Biologique ✓

temporary variations in pigment in prodigiosus:

La aut. My. in pigm prod.

Brown, F.M. + H.M. Heffron, Science 49:198-200 (1929). Mendelism among bacteria?

yellow "bacillus"

*B. lutyus* (Brown)

G+ young cultures


"sph. refractile bodies" in old cult.

- a) yeast like organism associated "both while annoying at the time have been found to be merely phases in the life history..."  
b) loss of color.

a) filament formation in old cultures

- b) sudden toxin: e.g. Pb. : 8 rods fuse into a mass, staining intensely  $\bar{c}$  fusulin "symplesm."

Mass ~~then~~ divides into 4 sph. non staining bodies. Each of these  $\rightarrow$  tetrad  $\rightarrow$  16 "cocci" On transfer to new medium, cocci divide and  $\rightarrow$  rod form.



by white strains, old cultures, or symplesm formation  $\rightarrow$

both yellow + white colonies. Each, best time is daily transfer.

Single cell isolations of each made 9 2 wks. After 11 transfers, "substrains that showed no change of color from the 1st. single cell isolations were taken and considered to be pure strains of that color". Serologically identical.

Both cultures were mixed. On transfer, almost entirely white. (Fountain Valley School, Colorado Springs, Colo);

isolated symplasmas from mixed cultures on Pb-untreated agar slant  
grown in broth  
and plated out.  $\rightarrow 362:138 \text{ w:y. } (8:3)$

$\rightarrow w+y \quad w > y.$


In one instance  $\rightarrow$  all white.

V. 144p: single cell isolation after mixed culture???

1. Pure strains stable
2. Mixed cultures  $\rightarrow$  unstable white colonies.

Assume that there is a diploid segregation in  $F_1$   $\hat{=}$  same name.

1) Should have studied the progeny for variance.



Kowen + Imich J Bact 44:551- (1942) A test for sexual fusion in bacteria

1. yellow + wh. strains of *Phytonomasteter*.

a) Look for heterozygous colonies, septing colonies from mixture. Also mated by R, S. No recombination found.



4/20/22 mixed colonies found. Stable on septating (this is not unexpected in view of a small % of sticking together).

2. Recombination of characters (haploid)

509 - large, smooth, mucoid, wh. 400 sm, rough, non muc, yellow.

Mutations observed in parentals, as frequently, as in mixed culture.

↓  
app. recombinations.



Data not analysed.  
> 200 000 colonies examined.  
no literature.

See J Bact 33.

Erkay, L + W. Hesselbrock, J Bact 49: 235 - . 1945. Some observations  
on the filterability of *M. tularensis*.

Filter ca 300-350 m $\mu$  demonstrated by infectivity + subculture.

Immunelski A J Biol ~~Sci~~ 49:1-5 (1945) On the structure of anaerobic  
bar-

Hollande, Arch Protistenk. 83:465-608 (1934) Contrib. à l'étude  
cytologique des microbes (Eucyres...)

Dejeu, L. J. *Bact* 50: 441-458. Morphology + Nature of the  
Pleuropneumonia group of organisms.



(R)

Altman - Weber, E., et al J Bact 50: 291-5 (1945) The effect of  
incompletely inh. conc. of penicillin on E coli. Rept Lab; Jewish Hosp.  
Brooklyn, N.Y.

Nutrient broth:

75 units/ml Bact → "bipolar" leptothecoids?

Ab. at 300/ml

at 100, mycelium

150 "zygospores"  
200 early small cells.

Ade, P.A. J Bact 51(6): 699-701 (1946) Mutation in certain  
phytopathogenic bacteria induced by acenaphthene.

Pl Path  
98 Baskely

*Phytophthora michiganensis* + *Erwinia carotovora*

acenaphthene saturated nutrient broth. 2 vols 28°.

by *P. mich* "a sudden + complete mutation" → only a wh. shiny  
smooth type of colony. Neither intermediate nor typical forms  
were found after a certain time.

*E. carot* → several types - peid. grayish compact flat colony

Ramchandani, J. C. Ann. Bot. 44: 975-987 (1930) *Saltans*  
in *Bacteria*. III *B. violaceus*

color variations.

Seib. 40: 2, 43: 579.

→ wh. mutant + reversion.

Hosh. EC PRS 389:468 (1917). Morph St. in the life history  
of bacteria.

Breeding? (Centricity.)

Stewart FH J Hyg 27:379-95 (1928) The life cycle of *pasturia*,  
alternate asexual and autogamous phases.

Rosen HR Mycologia 20: 251-75 (1928) Variations within a  
bacterial sp - I Morphologic Variations Ark.

"Gurney-Dixon, S. "The transmutation of bacteria" 1919.

"B. mesentericus?" Tiny particles attached to flagellae were  
seen.  
interpreted as germinis.


No direct evidence of vitality. Filtrates did show culture.

∴

not relevant.

Beebe, J. H. J. Part 42: 193-223 (1941) The morphology and  
cytology of *Myxococcus xanthus*. m.s.  
Univ. Ariz. Tucson. (R)

Describes nuclei  $\bar{c}$  2 chromosomes and autogamous  
fission before sporulation. Meiosis not observed.



(Seems OK.) But a myxobacterium !!

Nyberg, C. Acta Soc Med Fennicae 12: 1-18 (1930)  
des *Bacillus myxoides*.

Zur Biologie



Broadhurst, J. & J. Bach 27: 48 (1934) SAB.  
*Zygote pheres in bacteria.*

[Find "Lommel" 1926; Wygodtshichoff + Mamulova 1930  
Roux + Yersin 1890.]

# Classification of Literature.

## 1. Growth

- a general
- b growth factors
- c " " - analogues
- d antibiotics
- e regeneration - see sp. organs
- f genetics.

## 2. Genetics

- a transmission
- b gene acting as gene; induced mutation.
- c action
- d biochemical, in microorganisms
- e other, "
- f adaptive enzymes.

JPB  
JCP  
JLEM  
JEM  
JCol Res  
J.O.Ch.  
J.Ph.Ch.  
Faraday Soc.  
J. Russ.  
J. Hyg.  
J. Chem Soc.  
Exp Phys & Th.

Enzymologia  
Advances

Biol Rev.  
Zool Rev.  
Q.R. Biol

"Flux" in "specific" proteins not established.

Is order specific ???

Is order maintained in derivat.??

Are <sup>sp.</sup> proteins made by enzymes??

Spont. reactivation.

"Specificity" - <sup>+ substrate:</sup> enzymatic -  
immunologic  
[ genetic ]  
how else.

Boils J 39(5):/ 1945.

$$B \xrightarrow{m} B/r$$

$$r \xrightarrow{m} r'$$

$$r' \propto B, B/r$$

$$Br \xrightarrow{m} Brd.$$

$$\alpha \not\propto Brd$$

$$r' \propto Brd.$$

$\therefore$  this mutational resistance is specific.

$$Bd, \xrightarrow{m} Bd, r'$$

$$r', r \not\propto Bd, r'$$

$$\alpha \propto Bd, r'$$

!!!

$$Bd, r' \xrightarrow{m} Bd, r' \alpha$$

resistant to  $\alpha, r, r'$ .

The mutant viruses are all active in original host!

$$B \xrightarrow{m} Bd_1.$$

$$B \xrightarrow{m} Bd_2. \text{ small colony mutant on nutrient agar!}$$

$$T1 = \alpha$$

$$T2 = r$$

Some Aspects of the Nitrogen Metabolism of a Lysogenic  
Strain of Bacillus megaterium

The total nitrogen of the infected and uninfected cells was determined by the semi-micro Kjeldahl technique. The uninfected cells were found to contain a larger amount of total nitrogen than the infected cells. It was found that the desoxyribonucleic acid (DNA) content of the infected cells was slightly higher than that of uninfected cells. The presence of the virus in lysogenic cells in the immature form is believed to be the explanation for the slightly larger amount of DNA in the infected cells.

The technique of Feldman and Gunsalus was used to study the activity of the transaminases of B. megaterium. Pyridoxal-PO<sub>4</sub> was required as a coenzyme and a number of amino acids could serve as amino donors.

The effect of sodium azide, sodium fluoride and iodoacetate on growth and virus production was studied. NaF had little or no effect in the concentrations used. Sodium azide and iodoacetate depressed growth and virus formation. The inclusion of ATP in the medium, along with the inhibitor, produced inconclusive results.

Studies with N<sup>15</sup>-ammonium carbonate showed that after a 30 min. exposure the amount of N<sup>15</sup> taken up by both strains was the same. A study of N<sup>15</sup> distribution in amino acids, purines and pyrimidines was done also.

A complex amino acid medium was developed; it supported better growth of the lysogenic strain than nutrient agar, the amount of virus produced was significantly less. The addition of asparagine or adenine to the amino acid medium increased virus formation.

The two strains of B. megaterium were grown on synthetic media containing purines, pyrimidines and nucleotides as the sole sources of nitrogen. The uninfected cells showed good growth on these media, but the growth of the lysogenic strain was only fair. The lag phase could be shortened appreciably with larger inocula, i. e., direct transfer. Little or no ammonia was liberated, and there was little change in pH over a 48 hour growth period. Attempts to isolate and identify intermediate products of metabolism by chemical methods, paper chromatography and UV irradiation were unsuccessful.

References

- Feldman, L. I., and Gunsalus, I. C. 1950 The occurrence of a wide variety of transaminases in bacteria. J. Biol. Chem., 187, 821,830.
- Lwoff, A., and Gutman, A. 1950 Recherches sur un Bacillus megaterium lysogene. Ann. inst. Pasteur, 78, 711-739.

4/52

Rhoades *ijap*.  
"Hunt's ... 1950"

①  $I_j i_j$  shows no effect.

② Give  $i_j i_j$  segregants

which are stupid. ③ Stupid segregants  $\times I_j \rightarrow$  stupid

$I_j i_j F_2$ . ④ Stupid  $F_2 \times \bar{I}_j \bar{i}_j \rightarrow$  stupid  $I_j I_j$  (genetic marker).

1.  $\therefore$  Pleistid abnormality is inherited at least two generations in presence of  $I_j$ . Further selfing of  $I_j I_j$   $\rightarrow$   $\pi$ .

⑤ Virus is brought in from  $i_j I_j$  stock. This virus has no effect in presence of  $I_j$  but can be propagated in presence of  $\bar{I}_j$ .

See Jenkins MT J Her. 15: 467-472.

Notes that green  $\bar{I}_j i_j$  plants show "conditioning"

In summer-grown plants,  $F_1$  plants are pure green. In out-of-season plants (req. 4 mos for maturity) white-stripping is seen: intracellular competition. Lectus modifies *ijap* pleistids in same sense as others.

Other genes do not behave in the same way. (1948)

Dojap.  $y/j$  are striped.

$y \sigma \times Ij \rightarrow$  normal  $F_1$ .  $y \text{♀} \times Ij \rightarrow$  white and striped  
as well as green  $F_1$ .  
 $\nearrow$   
 $F_2$   
3:1  $Ij:yj$

$yj$  plastids are smaller as well as chlorotic. "Both types of plastids were found in certain green cells."

Striped  $F_1$  ( $yj \text{♀} \times Ij \sigma$ )  $\times$  unstriped  $Ij$   $\rightarrow$  plants  $1/2$  should be  $Ij Ij$ .

Occasionally all progeny of a backcross ear (white sector) were white seedling, though  $1/2$  were  $Ij Ij$ . Concludes that mutant plastids retain their individuality.

(Persistence of striping in  $Ij Ij$  striped plants?)

Later, glossy-1 was used to mark  $Ij$  to prove homozygous condition. Normal sized plastids were paler in cells adjacent to white tissue. Also proportion of white offspring less than expected from proportion of maternal white tissue. Direct effect of  $yj$  on cytoplasm supposed: indecisive whether the permanent changes are in cytoplasm or plastid: plastid or plasmagene mutation? Segregation as ear of Rimm case is best evidence for cyto factor



- x Aschner, M. 1946 The symbiosis of Lucanosipoda aegyptia Mcq. Bull. Soc. Fouad Entom., 30, 1-6.
- x Borghese, E. 1947 Il ciclo del batterio simbiote di Blattella germanica (L.). Mycopathologia 4. (18 pp)
- Borghese, E. 1948 Osservazioni sul batterio simbiote della Blattella germanica L. Monitore Zool. Italiano 56 (Suppl.), 252-253.
- x Bramstedt, F. 1948 Über die verdauungsphysiologie der aphiden. Zeit. f. Naturforschung, 3b, 14-24.
- ✓ Buchner, P. 1947 Symbiosis in Oliarius. Nature, 160, 264. *cf. Mahdihassan 1947*
- Buchner, P. 1949 Symbiose der Tiere mit pflanzlichen Mikroorganismen. Sammlung Götschen, 1128, 128 pp.
- Csáky, T. Z., and Tóth, L. 1948 Enzymatic breakdown of nitrogen compounds by the nitrogen fixing bacteria of insects. Experientia, 4: 4 pp.
- ✓ Hier, H. T. 1947 Intracellular bacteroids in the cockroach (Periplaneta americana Linn.). J. Bact., 53, 173-189.
- ✓ Glaser, R. W. 1946 The intracellular bacteria of the cockroach in relation to symbiosis. J. Parasitol., 32, 483-489.
- Goetsch, W. 1946 Darm-Symbionten als Eiweißquelle und Vitamin-spender. Österreichische Zool. Zeit., 1, 58-86.
- Goetsch, W. 1946 Vitamin "T" ein neuartiger Wirkstoff. Österreichische Zoologische Zeitschrift, 1, 49-57.
- x Gubler, H. U. 1947 Versuche zur Zucht intracellulärer Insekten-symbionten. Inaugural-Dissertation, Univ. Zürich. 34 pp.
- x Keller, H. 1950 Die Kulture der intrazellulären Symbionten von Periplaneta orientalis. Z. f. Naturforschung, 5b, 269-272.
- x Koch, A. 1948 Begriff und Bedeutung der Symbiose. Mikrokosmos, 38, 4 pp.
- ✓ Koch, A. 1948 Wege und Ziele der experimentellen Symbiosforschung. Naturwiss. Rundschau, No. 4, October, 1948, pp 166-171.
- x Koch, A. 1949 Die Bakteriensymbiose der Küchenschaben. Mikrokosmos, 38, 6 pp.
- Koch, A. 1950 Fünfzig Jahre Erforschung der Insektensymbiosen. Die Naturwissenschaften, 37, 313-317.
- Mahdihassan, S. 1946 The micro-organisms in Melophagus ovinus. Current Sci., 15, 166-167.

↓ Mahdihassan, S. 1946 Bacterial origin of some insect pigments. Nature, 158, 58-59.

• Mahdihassan, S. 1947 Cicadella viridis, Its symbiotes and their function. Curr. Sci., 16, 58-59.

✓ Mahdihassan, S. 1947 Specificity of bacterial symbiosis in Aphrophorinae. Proc. Ind. Acad. Sci. 25, 155-162.

Mahdihassan, S. 1947 Bacterial symbiosis in a Margarodes sp. Current Science, 16, 379-380.

\* Mahdihassan, S. 1947 Bacterial symbiosis in Aphis rumicis. Acta Entomologica Musei Nationalis Pragae, 25, 123-126.

af Mahdihassan, S. 1948 Bacterial origin of muscle pigment in a Cicada. Rev. de Entomologia, 19, 585-586.

\* Peklo, J. 1946 Symbiosis of azotobacter with insects. Nature, 158, 795-796.

\* Rau, A. 1943 Symbiose und Symbiontenerwerb bei den membraciden (Homoptera-Cicadina) Zeit. Morphol. Okol. Tiere, 39, 369-522.

, Schanderl, H. 1949 Unsere derzeitigen Kenntnisse über die Hefen. Biologia Generalis, 19, 130-162.

✓ Schanderl, H., Lauf, G., Becker, H. 1949 Studien über die Mycetom- und Larvymbionten der Aphiden. Zeit. f. Naturforschung, 4b, 50-53.

\ Tóth, L. 1946 The biological fixation of atmospheric nitrogen. Monographs in Nat. Sci., No. V, Hungarian Museum of Nat. Sci. 116 pp.

• Tóth, L. 1950 Protein metabolism and nitrogen fixation by means of micro-organisms living in symbiosis with insects. Eighth International Congress of Entomology, 303-306.